The Murderer of the Missoula Floods • G202 Winter 2015, SE Center

By: Velah Kimble, Julianna Jones, Luke Lorenz, Kvernland Vance

When to Worry

- Naturally occurring
- Only an issue when trapped in home

Radon's In The House

- Enters through water and small cracks
- Gets trapped and concentrated




Figure 1 Radon gas entering a house through cracks, pipes, drains, and other small openings,

http://ohiohomeanswers.com/image_store/uploads/8/8/2/6/0/ar1285609237 06288.jpg

Uranium to Radon— Dude!

- Uranium in soil
- Goes through multiple transformations
- Ends up as radon




Figure 2. Diagram showing Uranium to Radon decay chain, http://www.mdpi.com/energies/energies-07-

08483/article_deploy/html/images/energies-07-

- If experiencing symptoms such as:
- Pressure in chest
- Trouble breathing
- Dry, persistent cough
- Feeling of drowning
- If symptoms persist or get worse
- If house is old or spend lots of time basement
- If radon levels in house 4pCi/L or higher
- pCi/L = picocuries per liter
- Used to measure radiation
- Curie (Ci) = 1 gram pure radium
- Pico (p) = 1×10^{-12}
- 1pCi/L = 2.5 cigarettes a day

Breasts vs. Lungs

- Third biggest killer in US behind Lymphoma and Leukemia
- Society is very focused on other forms of cancer, such as prostate and breast with lung cancers being left out of this movement.
- Gets under \$2,000 Federal funding/year
- Breast cancer + prostate cancer = 1/3 number of radon cancer deaths/year combined
- Each get over \$10,000 Federal funding/year alone

Figure 3. When-to-

http://www.portlan

dradonmitigation.c

om/core/images/ra

mitigation/radon-

information/radon

-levels/radon-

graphic.png

act chart,




Figure 5. Amount of Federal funding per cancer per year,

https://s-media-cache-


ak0.pinimg.com/236x/ff/e2/d5/ffe2d534a02f4e196a04da3cd4395a68.jpg

4.0 The EPA recommended 2.7 World Health Organization recommended radon action level. radon action level. 3.0 5.0 pCi/L 2.0 The amount of radiation in 4.0pCi/L IpCi/L of radon is equal to 2.5 of radon is equivalent to that in 100 cigarettes a day. The EPA states chest X-rays. (Most hospitals only allow radon is the #2 cause of lung people to have 4 chest x-rays or cancer - only smoking is #1. .64 pCi/L each year.)

Cancerous effects of Radon

- Over 20,000 USA deaths/year due to radon gas
- Second leading cause of lung cancer next to smoking
- IS RADIATION!
- Alpha particle
- Have to breath in to get sick
- EVERYONE has to breathe

Figure 4. Radon Lung Cancer, http://ohiohomear wers.com/image_s re/uploads/3/7/1/9/ /ar12856088645917



I Will Survive

- Survivable with early detection
- Tumors can be removed
- Smoking increases risk DRASTICALLY
- If smoking, stopping can save you life
- Live in older building or basement? GET A CHECK-UP AND X-RAY!




Figure 6. Quote from a lung cancer survivor,